


TAR SPOT RATINGS



Tar spot is a foliar disease that can affect the yield output of corn hybrids. Below you will find tar spot ratings for our 2026 corn product lineup. We have also included frequently asked questions about this disease. Please refer to the Burrus Harvest Report, Product Selection Guide or our website at burrusseed.com for more information on our products. As always, you can also contact your Burrus Representative or Field Agronomist with any additional questions.

PRODUCT	RATING
Burrus 7K244 PCE	7
Power Plus® 1K12AM*	7
Power Plus® 1U41AM*	8
Burrus 1R62 PCE	6
Burrus 2A13 PCE	7
Power Plus® 3G31AM*	6
Burrus 4W42 PCE	7
Power Plus® 4V73AM*	7
Burrus 5C11 PCE	6
Power Plus® 5J21AM*	7
Burrus 6L42 PCE	7
Burrus 6M12 AA	6
Burrus 6N54 VT2P	5
Burrus 6Z72 TRE	5
Burrus 7B14 PCE	6
Burrus 7F33 VT2P	5
Burrus 7Y23 TRE	6
Burrus 8A12 VT2P	5
Burrus 8K34 VT2P	5
Burrus 8C51 PCE	7

PRODUCT	RATING
Power Plus® 7K299 VE*	7
Power Plus® 1K18 Q*	7
Power Plus® 1R47 VE*	6
Power Plus® 2A17 VE*	7
Power Plus® 2J67 Q*	6
Power Plus® 3H48 VE*	7
Power Plus® 4P27 VE*	7
Power Plus® 5M76 VE*	7
Power Plus® 5W86 VE*	8
Power Plus® 6G88 VE*	6
Burrus 6U77 SSP	5
Burrus 7T27 SSP	5
Power Plus® 7R68 VE*	6
Power Plus® 4W40*	7
Power Plus® 5W20*	8
Burrus 6L40	7
Burrus 7F30	5

RATINGS: 10 = BEST; 1 = POOREST

WHAT IS TAR SPOT?

Tar spot is a foliar disease native to Latin America that made its way to the Midwest in 2015. It has been most problematic in 2018 and 2021 due to weather conditions favorable for its development. The disease dramatically expanded its geographic reach in 2021. Tar spot produces dark structures called ascomata that develop on corn leaves. These fruiting structures are impregnated into the leaf and cannot be rubbed or scraped off without damaging the plant. Each ascomatum produces spores that can be windblown long distances. Additionally, these structures can overwinter on corn residue, releasing spores the following season. While differences in hybrid susceptibility to tar spot do exist, there is currently no known commercially available germplasm that is completely resistant.

HOW DID WE DETERMINE TAR SPOT RATINGS?

Our ratings were compiled through opportunistic observation over multiple locations and in some instances, multiple years. This allows for a fairly accurate assessment of tar spot susceptibility in our lineup; however, ratings can be influenced on the initial presence of tar spot and the environmental conditions sustaining its development. Hybrid ratings may fluctuate over time as we continue to monitor susceptibility; however, there is no known process currently to inoculate trials with tar spot to facilitate uniform disease pressure and timing.